

Local epidemiology in HDV

Marina Berenguer & Helena
Hernandez

Search for patients



HBV PATIENTS



Anti-HDV

Collect Variables



PARAMETERS

- Demographics
- Epidemiologic
- Clinical
- Serological
- Virological


Analyse data



605 HBV patients



70 HDV patients (11.57%)



62.9%

Median age: 52 yrs

Country of origin



67.1%

(Easteern Europe: 24.3%)

Transmission

45.7% unknown

20.7% nosocomial

14% IVDU



Comorbidities



Diabetic: 8.6%



Overweight/obese (BMI>25.5): 55.5%



Alcohol (past/present): 34.1%




Viral coinfection: HCV 11.4%; HIV 10%

Comorbidities


Transmission
Nosocomial
intrafamily



Overweight/
Obesity: 35%



Alcohol: 49%



Transmission
IVDU
Nosocomial


Overweight/
Obesity: 80%




Alcohol: 88%



Therapies



0 long-term responses



77.1%



90.7% men cirrhotic

Risk factors:


- Older age
- Diabetes Mellitus
- Lack of antiviral therapy

17.4%



Risk factors:


- Older age
- Alcohol




59.3%

Differences overtime

PRE-2000



POST-2000



Conclusion

Look for it and you will find it

Evangelii segons Sant Mateu 7:7

